

INITRO

Nitroxylnil 34% w/v Injection



COMPOSITION :

Each ml contains :

Nitroxylnil B.P	340 mg
Water for Injections B.P	q.s.

INDICATION :

INITRO 34% Injection is effective against:

- Liver fluke infestations caused by Fasciola hepatica and Fasciolagigantica.
- Gastrointestinal parasitism caused by Haemonchus, Oesophagostomum and Bunostomum in Cattle, Sheep and Goats.
- Oestrusovis in Sheep and Camels.
- Hookworms in Dogs.

CONTRAINDICATION :

Do not use in animals with known hypersensitivity to the active ingredient.
Do not use in animals producing milk for human consumption. Do not exceed stated dose.

DOSAGE AND ADMINISTRATION :

For Subcutaneous Cattle, Sheep, Goats, Camels : 1ml per 35 kg bodyweight by Subcutaneous Injection. This dosage may be increased to 1.25 ml per 35 kg bodyweight in case of acute fascioliasis (immature flukes).
Dogs : 0.30 ml per 10 kg bodyweight by Subcutaneous Injection.

WITHDRAWAL PERIOD :

Withdrawal times :

For meat :

Cattle : 60 days.

Sheep : 49 days.

STORAGE :

Store at a temperature not exceeding 30°C.
Protect from light.



Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com