

AMPROCIN 20%

AMPROLIUM WATER SOLUBLE POWDER 20% W/W FOR ORAL ADMINISTRATION



COMPOSITION :

Each gm contains :

Amprolium hydrochloride	B.P.	20% w/w
Excipients		q.s.

INDICATIONS :

Amprocin 20% is indicated for the treatment & control of caecal and intestinal coccidiosis in Poultry caused by *Eimeria tenella*, *E. maxima*, *E. brunetti*, *E. mitis* & *E. acervulina*.

DOSAGE AND ADMINISTRATION :

Amprocin 20% can be administered through drinking water.

Poultry :

Mild Outbreak : 5 g of Amprocin 20% per 8 lt. of drinking water for 5 - 7 days.

Severe Outbreak : 5 g of Amprocin 20% per 4 lt. of drinking water for 5 - 7 days.

Calves, Goats & Sheep :

Prevention : 5 mg / kg bodyweight or 1 g of Amprocin 20% per 40 kg bodyweight, through drinking water or milk for 21 days.

Treatment : 10 mg / kg bodyweight or 1 g of Amprocin 20% per 20 kg for 5 days.

CONTRAINDICATION :

The use of Amprolium is prohibited from a laying age onwards. Do not administer to Poultry whose eggs are intended for human consumption, or to animals with impaired liver and/or renal functions. Do not administer to turkeys before the age of 8 to 10 weeks.

WITHDRAWAL PERIOD :

Animals must not be slaughtered for human consumption during treatment.

The withdrawal periods after the last treatment are :

Meat

Poultry : 3 days

Calves, Goats & Sheep : 3 days

PACK SIZE :

100 gram, 500 gram Sachet & 1 kg of Jar.

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

