



BUPACIN

BUPARVAQUONE INJECTION 50 MG/ML



COMPOSITION :

Each ml contains :

Buparvaquone	50 mg
Sorbitan mono oleate B.P.	100 mg
Oil base	q.s.

INDICATIONS :

It is indicated for theileriosis (East coast fever, Corridor disease, Zimbabwean Theileriosis, Tropical Theileriosis) caused by intracellular protozoan parasites *Theileria parva*, *Theileria parva bovis*, *Theileria parva lawrencei*, *T.annulata*, *T.mutans* and *T.orientalis sergenti* in Cattle. It is active against both the schizont and piroplasm stages of *Theileria* spp., and may be used during the incubation period of the disease, or when clinical signs are apparent.

CONTRAINDICATIONS :

Due to the inhibiting effects of theileriosis on the immune system, Vaccination should be delayed until the animal has recovered from theileriosis.

DOSAGE AND ADMINISTRATION :

A single dose of 1 ml BUPACIN Injection per 20 kg bodyweight. (2.5 mg buparvaquone per kg bodyweight). In severe cases the treatment may be repeated within 48-72 hours. Not more than 10 ml should be injected into a single site. Successive Injection should be administered at different sites.

ADMINISTRATION :

Intramuscular Injection into the neck muscles.

WITHDRAWAL PERIOD :

Milk : 2 days

Meat : 42 days

PACK SIZE : 20 ml, 50 ml, 100 ml Vial

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

