



BUPACIN PLUS



BUPARVAQUONE WITH FUROSEMIDE INJECTION

COMPOSITION :

Each ml contains :

Buparvaquone		50 mg
Furosemide	B.P.	55 mg
Sorbitan mono oleate	B.P.	100 mg
(As preservative)		
Oil base		q.s.

INDICATIONS :

Indicated for the treatment of theileriosis (East Coast Fever), particularly in advanced cases associated with pulmonary oedema.

Buparvaquone is active against the schizont (lymphoid cell) and piroplasm (erythrocytic) stages of Theileria spp., and suppresses pre-schizont stages during the incubation period. It acts by inhibiting the parasite's mitochondrial electron transport (respiratory) system.

Furosemide is a diuretic that helps reduce pulmonary oedema.

DOSAGE AND ADMINISTRATION :

Cattle & Sheep :

Injection by the Intramuscular route into the neck muscles at the rate of 1 ml / 20 kg (2.5 mg per kg of Buparvaquone and 2.75 mg/kg of Furosemide) Repeat after 48 hours.

In cases of exceptionally severe infections with pulmonary oedema, further treatment at half the dosage rate may be required at 24 hour interval.

WITHDRAWAL PERIOD :

Milk for human consumption should not be taken from animals treated with BUPACIN PLUS until atleast 48 hours after treatment. Milk for animals treated with BUPACIN PLUS is safe for consumption by Calves. Animals should not be slaughtered for human consumption until at least 42 days after treatment with BUPACIN PLUS.

PACK SIZE :

20 ml, 50 ml, 100 ml Vial

Veterinary Medicines - Injectables

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

